Forces, Energy and Power

- force: acts on body
- energy: work done on body
- or

capacity to do work

• power: work done per time

Newton's three laws of motion: [¶]

- 1. → An object continues in a state of rest or in a state of motion at a constant speed along a straight line, unless compelled to change that state by net force. ¶
- 2. → When a net external force, F, acts on an object of mass, m, the acceleration, a, that results is proportional to the force and has a magnitude that is inversely proportional to the mass. The direction of the acceleration is the same as the direction of the force.
- 3. → Whenever one body exerts a force on a second body, the second body exerts an oppositely directed force of equal magnitude on the first body. ¶

-

Example: 2 Guys Pushing a Car



F proportional to:

mass *m* acceleration *a*

$$F = m x a$$

Work *W* proportional to: force *F*

distance D

$$W = F \times D$$

Power *P* proportional

toork W inverse of time t

$$P = W/t$$

SIO15: Chapter 2: Forces, Energy and Power

Example: 2 Guys Pushing a Car

Power *P* proportional toork *W* inverse of time *t*

$$P = W/t$$



If same work is done in less time, then the process required more power. E.g. push a car in 10 min over distance D requires more power than doing this in 20 min.

Gravitational Force

$$F = m x a$$



attraction of bodies due to their mass

For bodies that are small compared to Earth and are close to Earth's surface, compared to its radius (6371 km), we can assume that g is constant.

gravitational acceleration g

$$Fg = m x g$$



Gravitational Force and Potential Energy

Example: Isaak Newton's Apple



When the apple hangs on the tree, it has potential energy that can be used up and transferred into another type of energy.

SIO15: Chapter 2: Forces, Energy and Power

$$Fg = m x g$$

Epot proportional to:

mass *m* gravitational force *Fg* height *h*

$$Epot = Fg \ x \ h$$
$$= m \ x \ g \ x \ h$$

Gravitational Force, Potential Energy and Kinetic Energy

Example: Isaak Newton's Apple



When the apple falls, it loses potential energy and gains kinetic energy. When it reached the ground, it lost all its potential energy. Just before it hits the ground, it has only kinetic energy. When it hits the ground, this energy is used up to deform the apple.

Exchange of Potential and Kinetic Energy

energy can be transferred from one type to another

e.g. when a mass slides down a slope



 $Epot = m x g x h \qquad Ekin = 1/2 m x v^2$

This example ignores friction between the mass and the slope. If friction is included, some of the potential energy is lost to heat for the moving mass to overcome the friction. SIO15: Chapter 2: Forces, Energy and Power

Elastic Potential Energy in a Loaded Spring

relaxed state loaded state Spring can be loaded only to Χŋ Хı a certain point $X = X_1 - X_0$ before it no

When you pull on a spring that was initially relaxed, you use a force to load it. This force is proportional to the distance over which you pull, and the spring constant, k,that describes the stiffness of the spring (or its resistance to the pull).

$$Fg = k \times x$$

with: *k*: spring constant *x*: distance pulled

Epot =
$$1/2 k \times x^2$$

Though the math is more complex, the same principle applies to elastic media that are bent without breaking them (e.g. a wooden stick, lithospheric plate). When the medium breaks, Epot is released.

SIO15: Chapter 2: Forces, Energy and Power

longer behaves

elastically and

breaks and.

Elastic Potential Energy in a Lithospheric Plate

In earthquake science, we use the words stress and strain instead of force and distance pulled (deformation). But the analogy to a loaded spring is obvious.

force <-> stress

deformation <-> strain

strain is a result of stress

The potential energy accumulated during the loading is released in an earthquake. This energy is used when you feel the shaking (Ekin) and structures are

SIGAMAGE Procest, exercity and Power



Properties of a Material



SIO15: Chapter 2: Forces, Energy and Power

Fault

Gravitation

attraction of bodies due to their

mass attracting force: gravitational force *Fg*

If two bodies have about the same mass (e.g. Earth and Moon) then the gravitational force is more complicated:

Newton's Apple



m: mass of attracted body *g*: Earth's gravitational acceleration

Moon MM

$$Fg = m \times g$$

M_E: Earth's Mass *M_M*: Moon's Mass *d*: Earth-Moon distance *G*: Gravitational constant

$$Fg = \frac{G \times M_E \times M_M}{d^2}$$



Gravitation

attraction of bodies due to their

mass attracting force: gravitational force *Fg*

If two bodies have about the same mass (e.g. Earth and Moon) then the gravitational force is more complicated:

If we consider a force with respect to Earth, then we can simplify by defining the gravitational acceleration, g, as:

Now we have an equation very similar to that on page 1, but g depends on the distance, d, between the two

SIO15: Chapter 2: Forces, Energy and Power

M_E: Earth's Mass *M_M*: Moon's Mass *d*: Earth-Moon distance *G*: Gravitational constant

$$Fg = \frac{G \times M_E \times M_M}{d^2}$$

$$g = \frac{G \times M_E}{d^2}$$

m: mass of attracted body *g*: Earth's gravitational acceleration

$$Fg = m x g$$

Gravitation



The Moon and Earth: Different Types of Energy



The Moon and Earth: Different Types of Energy

